

Drug Information Sheet("Kusuri-no-Shiori")_Draft

Internal

Published: 10/2015

The information on this sheet is based on approvals granted by the Japanese regulatory authority. Approval details may vary by country. Medicines have adverse reactions (risks) as well as efficacies (benefits). It is important to minimize adverse reactions and maximize efficacy. To obtain a better therapeutic response, patients should understand their medication and cooperate with the treatment.

Brand name: Neurovitan Tablets

Active ingredient: Octotiamine

Riboflavin

Pyridoxine hydrochloride

Cyanocobalamin

Dosage form: pale red tablet, diameter: 7.1 mm, thickness: 3.3 mm

Print on wrapping: (face) ノイロビタン, f536, (back) ノイロビタン, ビタミン B 群配合錠, company mark



Effects of this medicine

This medicine is a formulation of vitamins B₁, B₂, B₆ and B₁₂ to supplement these vitamins in short supply and restore balance.

It is usually used to supplement vitamins when they cannot be taken properly from food, or to treat neuralgia, myalgia/arthritis and peripheral neuritis/peripheral nerve palsy caused by vitamin deficiency or metabolic abnormality.

Before using this medicine, be sure to tell your doctor and pharmacist

- If you have previously experienced any allergic reactions (itch, rash, etc.) to any medicines.
- If you are pregnant or breastfeeding.
- If you are taking any other medicinal products. (Some medicines may interact to enhance or diminish medicinal effects. Beware of over-the-counter medicines and dietary supplements as well as other prescription medicines.)

Dosing schedule (How to take this medicine)

- Your dosing schedule prescribed by your doctor is << _____ to be written by a healthcare professional >>
- In general, for adults, take 1 to 3 tablets daily. The dosage may be adjusted according to the diseases, age or symptoms. Strictly follow the instructions.
- If you miss a dose, take a dose as soon as you remember. However, if it is almost time for the next dose, skip the missed dose and follow your regular dosing schedule. You should not take two doses at one time.
- If you accidentally take more than your prescribed dose, consult with your doctor or pharmacist.
- Do not stop taking this medicine unless your doctor instructs you to do so.

Precautions while taking this medicine

- Your urine becomes yellow because of vitamin B₂ formulated in this medicine, so it is nothing to worry about.

Possible adverse reactions to this medicine

The most commonly reported adverse reactions include abdominal distention, constipation, nausea, diarrhea and dizziness. If any of these symptoms occur, consult with your doctor or pharmacist.

The symptoms described below are rarely seen as initial symptoms of the adverse reactions indicated in brackets. If any of these symptoms occur, stop taking this medicine and see your doctor immediately.

No pertinent entries.

The above symptoms do not describe all the adverse reactions to this medicine. Consult with your doctor or pharmacist if you notice any symptoms of concern other than those listed above.

Storage conditions and other information

- Keep out of the reach of children. Store away from direct sunlight, heat and moisture.
- Discard the remainder. Do not store them.

For healthcare professional use only / /

For further information, talk to your doctor or pharmacist.